arXiv:1210.3280v2 [nucl-th] 20 Dec 2012

New Properties of High Momentum Distribution of Nucleons in Asymmetric Nuclei.

Misak M. Sargsian
Department of Physics, Florida International University, Miami, FL 33199 USA
(Dated: December 21, 2012)

Based on the recent experimental observations of the dominance of tensor interaction in the $\sim 250-600$ MeV/c momentum range of nucleons in nuclei, we suggest the existence of *two new properties* for high momentum distribution of nucleons in asymmetric nuclei. The *first* property is the approximate equality between proton and neutron high momentum distributions weighted by their relative fractions in the nucleus. The *second* property is the inverse proportionality of the strength of the high momentum distribution of protons and neutrons to the same relative fractions. Based on these two properties we modeled the high momentum distribution function for asymmetric nuclei and demonstrated that it describes reasonably well the high momentum characteristics of the ^3He nucleus. However the most surprising result we obtained is for nuclei with large A and asymmetry, for which we predict *a substantial per nucleon abundance* of high momentum protons relative to neutrons. For example, we predict that in the Gold the relative fraction of protons with momenta above $k_F \sim 260$ MeV/c is 50% more than that of neutrons. Such a situation may have many implications for different observations in nuclear physics ranging from medium modification of hadrons to properties of neutron stars.

One of the exciting recent results in studies of short-range properties of nuclei is the observation of the strong (by factor of 20) dominance of the *pn* short range correlations (SRCs), as compared to *pp* and *nn* correlations, for internal momentum range of $\sim 250-600$ MeV/c[1, 2]. This observation is understood[1, 3, 4] based on the dominance of the tensor forces in the NN interaction at this momentum range corresponding to average nucleon separations of ~ 1.1 Fm. The tensor interaction projects the NN SRC part of the wave function into the isosinglet - relative angular momentum, $L = 2$ state, almost identical to the D-wave component of the deuteron wave function. As a results *pp* and *nn* components of the NN SRC will be strongly suppressed since they are mainly defined by the central potential of the NN interaction with relative $L = 0$.

High Momentum Distribution of Nucleons in Nuclei and 2N SRCs: Due to the short range nature of NN interaction, the property of a A-nucleon bound state wave function, in which one of the nucleons has momentum p, such that $\frac{p^2}{2m_N} \gg |E_B|$ (binding energy), is defined mainly by the 2N interaction potential (V_{NN}) at relative momenta $k \sim p$, i.e.: $\Psi_A(p, p_2, p_3, , p_A) \sim \frac{V_{NN}(k)}{k^2} f(p_3, .p_A)$, where $\vec{p}_2 \approx -\vec{p} \approx -\vec{k}$ and $f(\cdots)$ is a smooth function of the momenta of non-correlated nucleons[5–7]. This leads to the approximate relation for nucleon momentum distribution at $p > k_F$, with k_F being the characteristic Fermi momentum of the nucleus:

$$n^A(p) \sim a_{NN}(A) \cdot n_{NN}(p) \tag{1}$$

where the momentum distribution of the nuclei is normalized as $\int n^A(p) d^3p = 1$ and the parameter $a_{NN}(A)$ can be interpreted as a probability of finding NN SRC in the given nucleus A. The function, $n_{NN}(p)$ is the momentum distribution of the nucleon in the NN SRC[5, 6, 8–10], where NN represents the combination of all possible isospin pairs.

If, following the above discussed dominance of the tensor interaction, we neglect the contributions from *pp* and *nn* SRCs one predicts that in the range of $\sim k_F - 600$ MeV/c:

$$n_{NN}(p) \approx n_{pn}(p) \approx n_{(d)}(p), \tag{2}$$

where $n_d(p)$ is the momentum distribution in the deuteron.

Let us now define the individual momentum distributions of proton ($n^A_p(p)$) and neutron($n^A_n(p)$) such that:

$$n^A(p) = \frac{Z}{A} n^A_p(p) + \frac{A-Z}{A} n^A_n(p) \tag{3}$$

and $\int n^A_{p/n}(p) d^3p = 1$. Then from Eqs.(1,2,3) it follows that if we neglect the contributions from *pp* and *nn* SRCs, then the high momentum distribution strengths of proton and neutron weighted by their fractions in the given nucleus will be defined by the same probability of being in the *pn* SRC in the relative D state. This brings us to the formulation of the *first property* of the high momentum distributions, in $\sim k_F - 600$ MeV/c region:

$$x_p \cdot n^A_p(p) \approx x_n \cdot n^A_n(p). \tag{4}$$

where $x_p = \frac{Z}{A}$ and $x_n = \frac{A-Z}{A}$.

Let us now consider a nucleus as a system consisting of two (proton and neutron) components. Because in our approximation the NN SRCs are identified only with *pn* correlations, with an increase of the nuclear asymmetry when the one component dominates the other, the per nucleon probability of lesser component to be in 2N SRC will be larger than that of the dominant component. This observation allows us to formulate the *second property* of the high momentum distribution: that *the probability of proton or neutron being in high momentum NN correlation is inverse proportional to their relative fractions (x_p or x_n) in the nucleus.* Based on this feature, the

TABLE I: Fractions of high momentum protons and neutrons in nuclei A

A	$P_p(\%)$	$P_n(\%)$	A	$P_p(\%)$	$P_n(\%)$
12	20	20	56	27	23
27	24	22	197	31	20

Eqs.(2) and (4) can be used to construct the high momentum ($k_F < p < 600$ MeV/c) distribution of protons and neutrons in nuclei in the following form:

$$n^A_{p/n}(p) \approx \frac{1}{2x_{p/n}} a_2(A,y)\cdot n_d(p) \tag{5}$$

where $a_{NN}(A) \approx a_{pn}(A,y) \equiv a_2(A,y)$ and the nuclear asymmetry parameter is defined as $y = |1-2x_p|$.

Fractions of High Momentum Component of Protons and Nuetrons in Asymmetric Nuclei: The most important prediction that follows from the *second property* is that the number of high momentum protons and neutrons became increasingly *unbalanced* with an increase of the nuclear asymmetry, y. To quantify this statement, using Eq.(5) one can calculate the fraction of the nucleons having momenta larger than k_F as:

$$P_{p/n}(A,y) \approx \frac{1}{2x_{p/n}} a_2(A,y)\int\limits_{k_F}^{\infty} n_d(p)d^3p, \tag{6}$$

where we extended the upper limit of integration to infinity assuming smaller overall contribution from the momentum range of ≥ 600 MeV/c. The results of the calculation of these fractions for medium to heavy nuclei, using the estimates of $a_2(A,y)$ from Ref.[10–14] and k_F from Ref.[15] are given in Table.I. As it follows from the table with the increase of the asymmetry the imbalance between the high momentum fractions of proton and neutron grow. For example, in the Gold the relative fraction of high momentum ($\geq k_F$) protons is 50% more than that of the neutrons.

Checking the High Momentum Features for ^3He Nucleus: One can check the validity of our observations for the ^3He nucleus using the exact calculations based on the realistic wave functions. For this we use the wave function based on the solution of Faddeev equation[16] using Argonne V18[17] NN potential parameterizations.

First, we check the validity of Eq.(4) which is presented in Fig.1. Here solid lines with labels "x_pn_p" and "x_np_n" in Fig.1(a) represent proton and neutron momentum distributions weighted by factors $\frac{2}{3}$ and $\frac{1}{3}$ respectively. As one can see the proton momentum distribution dominates the neutron momentum distribution at small momenta reflecting the fact that in the mean field the probability of finding proton is larger than neutron just because there are twice as much protons in ^3He. However at 300 MeV/c proton and neutron momentum distributions become close to each other and stay like this up to the internal momenta of 600MeV/c. This is the region dominated by tensor interaction.

As it follows from Fig.1(b) where the ratio of neutron to proton momentum distributions is calculated the approximation of Eq.(4) in the range of $300-600$ MeV/c is good on the level of 15%.

Next, we check the the validity of Eq.(5). For this we use the estimate of $a_2(^3He)$ from Ref.[14] and the deuteron momentum distribution n_d calculated using the same Argonne V18 NN potential[17]. The calculation based on Eq(5) is given by doted line in Fig.1(a). As it follows from this comparison the model of Eq.(5) works surprisingly well starting at 200 MeV/c up to the very large momenta $\sim 1GeV/c$. This reflects the fact that the center of mass motion effect as well as 3N SRCs are not dominant in the ^3He nucleus.



FIG. 1: (a) The momentum distributions of proton and neutron weighted by x_p and x_n respectively. The doted line represents the prediction for the momentum distribution according to Eq.(5). (b) The ratio of neutron to proton momentum distributions weighted by x_p and x_n respectively.

The final prediction which we check, is the one following from Eq.(6) according to which the neutron is more energetic in ^3He than the protons. For this we calculate the average kinetic energy and momentum of bound proton and neutron using the realistic wave function[16] which yields:

$$\begin{aligned} &< E^n_{kin} >= 18.4 \text{ Mev} \quad \text{and} \quad p^n_{rms} = 186 \text{ MeV/c} \\ &< E^p_{kin} >= 13.7 \text{ MeV} \quad \text{and} \quad p^p_{rms} = 160 \text{ MeV/c} \end{aligned} \tag{7}$$

Thus we conclude that all the observations concerning to the features of high momentum distribution in asymmetric nuclei is in reasonable agreement with the results following from the realistic ^3He wave function. This gives us some confidence that the same observations should be valid also for asymmetric nuclear with large A.

Restriction of the Model: In the the above made observations we neglected by the pp and nn SRCs as well as considered the pn correlations at rest neglecting their center of mass motion. However it is rather well established that in the momentum range of $k_F < p <$ 600 MeV/c the center of mass of the NN SRC moves in the mean field of the A-2 system with the momentum distribution that has a width proportional to k_F or more precisely to he RMS value of the single nucleon momentum[7, 18, 19]. This effect will clearly alter the predicted approximate equality in Eq.(4) since its validity is based on the assumption that the pn SRCs are in the D-state in the Lab frame of the nucleus (SRC at rest approximation, $|p_{p/n}| \approx k$, see discussion before Eq.(1)). However due to the mean field character of the center of mass motion it will affect equally the momentum distributions of proton and neutron in NN SRC. As a result one expects the validity of modified relation:

$$x_p^\gamma \cdot n_p^A(p) \approx x_n^\gamma \cdot n_n^A(p). \tag{8}$$

where $\gamma \equiv \gamma(k_F) \lesssim 1$, with γ decreasing with an increase of A (or k_F). The same γ factor will enter also in the high momentum part of the momentum distribution of the protons and neutrons

$$n_{p/n}^A(p) \approx \frac{1}{2x_{p/n}^\gamma} a_2(A,y) \cdot n_d(p), \tag{9}$$

which will diminish somewhat the imbalance between high momentum protons and neutrons presented in Table 1. One also expects a qualitatively similar effect due to pp and nn SRCs since they too will have an equal effect on the proton and neutron momentum distributions.

Implications of the Observed Features: The implications of above made observations are rather wide ranging, from the EMC effects to the proton properties in neutron stars. These observations also open up a new possibilities in studies of the high momentum component of nuclei with large A and asymmetry y.

-For example; combining three following observations: (i) nuclear medium modification of parton distribution functions (EMC effect) are proportional to the virtuality (high momentum) of the bound nucleon (see e.g. [9, 20, 21]); (ii) the high momentum protons dominate in nuclei with large A and y (this article) and; (iii) the deep inelastic structure function of proton dominates that of the neutron at $x_{Bjorken} \geq 0.3$ (see e.g. [22]) one can conclude that the EMC effects for large A asymmetric nuclei will be defined mainly by the proton component in the nucleus. This may explain[23] the large A part of the recently observed correlation between the strengths of the EMC and SRC effects[24, 25].

Prediction of the enhanced contribution of protons into the EMC effect indicates that in average the u-quarks will be more modified than d-quarks in large A asymmetric nuclei and the effect will grow with an increase of A. This may provide an alternative explanation of the NuTeV anomaly[26, 27]. The predicted effect also can be checked in parity violating deep inelastic scattering off the heavy nuclei.

- The discussed new features of the high momentum component will have significant impact on the proton distribution in neutron stars[14], where one expects $x_p \sim \frac{1}{8}$ and $y \sim \frac{7}{9}$. For example in Ref.[14] it was observed that starting at three nuclear saturation densities, protons will predominantly populate the high momentum part of the momentum distribution. This may have implication for several properties of neutron stars such as cooling through the direct URCA processes, superfluidity of protons, the magnetic field of the stars as well as the distribution of protons in the core of the massive neutron stars.

- Another possible implication of the high momentum protons in large A nuclei is that the volume occupied by protons will be smaller than the volume occupied by neutrons. This will create a new condition for formation of the neutron skin for large A asymmetirc nuclei.

Summary and Conclusions: Based on the dominance of the tensor forces in the NN system for momentum range of $\sim k_F - 600$MeV/c we observe a new equality relation between high momentum distributions of proton and neutron weighted by their fractions in the nuclei (Eq.4). Using this relation together with the fact of the dominance of pn SRCs in the high momentum part of the distribution function, we arrive at the second observation, according to which the strengths of the high momentum component of the proton and neutron are inverse proportional to the fractional contribution x_p and x_n. Based on these observations we constructed the high momentum distributions of protons and neutrons for asymmetric nuclei and using them estimated the overall fraction of nucleons being in the high momentum part of the momentum distribution.

Validity of our observations are confirmed by direct calculations using realistic wave function of ^3He.

We also observe that the effects due to center of mass motion of NN SRCs as well as contributions from pp and nn correlations will diminish the estimated imbalance between high momentum protons and neutrons in asymmetric nuclei. Thus we expect that our estimates represent the upper limit of such imbalance.

Our main conclusion for large A and asymmetry nuclei is that the high momentum distribution is dominated by protons. This dominance may have multitude of implications, some of which we discussed in the text.

It is worth noting that realistic nuclear structure calculations that can systematically incorporate short range correlations for asymmetric nuclei(see e.g. [28, 29]) are able to check the predictions of Eqs.(8,9) as well as evaluate the γ factor as a function of k_F. Experimentally, they can be checked in semi-inclusive high momentum transfer $(e, e'N)$ reactions on asymmetric nuclei at $x_{Bjorken} > 1$

and $|p_m^z| - \frac{q_0}{q_v}(E_m + \frac{p_m^2}{2M_{A-1}}) > k_F$, where p_m, E_m, q_0 and q_v are missing momentum, missing energy, transferred energy and transferred momentum in the reaction (see e.g. Ref.[30, 31] for details).

The author is thankful to Drs. J. Arrington, W.Boeglin, L. Frankfurt, O. Hen, D. Higinbotham, E. Piasetzky, A. Rios, M. Strikman and L. Weinstein for helpful comments and discussions. This work is supported by U.S. Department of Energy grant under contract DE-FG02-01ER41172.

---

[1] E. Piasetzky, M. Sargsian, L. Frankfurt, M. Strikman and J. W. Watson, Phys. Rev. Lett. **97**, 162504 (2006).
[2] R. Subedi *et al.*, Science **320**, 1476 (2008).
[3] M. M. Sargsian, T. V. Abrahamyan, M. I. Strikman and L. L. Frankfurt, Phys. Rev. C **71**, 044615 (2005).
[4] R. Schiavilla, R. B. Wiringa, S. C. Pieper and J. Carlson, Phys. Rev. Lett. **98**, 132501 (2007).
[5] L. L. Frankfurt and M. I. Strikman, Phys. Rept. **76**, 215 (1981).
[6] L. Frankfurt, M. Sargsian and M. Strikman, Int. J. Mod. Phys. A **23**, 2991 (2008).
[7] C. Ciofi degli Atti and S. Simula, Phys. Rev. C **53**, 1689 (1996).
[8] J. Arrington, D. W. Higinbotham, G. Rosner, M. Sargsian, Prog. Part. Nucl. Phys. **67**, 898 (2012).
[9] L. L. Frankfurt and M. I. Strikman, Phys. Rept. **160**, 235 (1988).
[10] L. L. Frankfurt, M. I. Strikman, D. B. Day and M. M. Sargsian, Phys. Rev. C **48**, 2451 (1993).
[11] K. S. Egiyan *et al.*, Phys. Rev. C **68**, 014313 (2003).
[12] K. S. Egiyan *et al.*, Phys. Rev. Lett. **96**, 082501 (2006).
[13] N. Fomin *et al.*, Phys. Rev. Lett. **108**, 092502 (2012).
[14] M. McGauley and M. M. Sargsian, arXiv:1102.3973 [nucl-th].
[15] E. J. Moniz, I. Sick, R. R. Whitney, J. R. Ficenec, R. D. Kephart and W. P. Trower, Phys. Rev. Lett. **26**, 445 (1971).
[16] A. Nogga, H. Kamada and W. Glockle, Nucl. Phys. **A689**, 357 (2001).
[17] R. B. Wiringa, V. G. J. Stoks and R. Schiavilla, Phys. Rev. C **51**, 38 (1995).
[18] M. Vanhalst, J. Ryckebusch and W. Cosyn, Phys. Rev. C **86**, 044619 (2012).
[19] A. Tang, J. W. Watson, J. L. S. Aclander, J. Alster, G. Asryan, Y. Averichev, D. Barton and V. Baturin *et al.*, Phys. Rev. Lett. **90**, 042301 (2003).
[20] W. Melnitchouk, M. Sargsian and M. I. Strikman, Z. Phys. A **359**, 99 (1997).
[21] M. M. Sargsian, J. Arrington, W. Bertozzi *et al.*, J. Phys. G **G29**, R1 (2003).
[22] J. Beringer et al. (Particle Data Group), Phys. Rev. D **86**, 010001 (2012).
[23] M. M. Sargsian,m arXiv:1209.2477 [nucl-th].
[24] L. B. Weinstein, E. Piasetzky, D. W. Higinbotham, J. Gomez, O. Hen and R. Shneor, Phys. Rev. Lett. **106**, 052301 (2011).
[25] O. Hen, E. Piasetzky and L. B. Weinstein, Phys. Rev. C **85**, 047301 (2012).
[26] G. P. Zeller *et al.* [NuTeV Collaboration], Phys. Rev. Lett. **88**, 091802 (2002) [Erratum-ibid. **90**, 239902 (2003)].
[27] I. C. Cloet, W. Bentz and A. W. Thomas, Phys. Rev. Lett. **102**, 252301 (2009).
[28] M. Alvioli, C. Ciofi degli Atti and H. Morita, Phys. Rev. Lett. **100**, 162503 (2008).
[29] A. Rios, A. Polls and W. H. Dickhoff, Phys. Rev. C **79**, 064308 (2009).
[30] L. L. Frankfurt, M. M. Sargsian and M. I. Strikman, Phys. Rev. C **56**, 1124 (1997).
[31] M. M. Sargsian, Int. J. Mod. Phys. E **10**, 405 (2001).